

**May 2025**  
**M.Sc.**  
**Fourth Semester**  
**CORE – 11**  
**PHYSICS**  
*Course Code: MPHC 4.11*  
(Computational Physics)

Total Mark: 70

Pass Mark: 28

Time: 3 hours

Answer five questions, taking one from each unit.

**UNIT-I**

1. (a) Using Newton-Raphson method, find the real root of the equation  $3x = \cos x + 1$ , correct up to four decimal places. 8
- (b) Find the relative error if the number  $x = 0.004997$  is  $3 \times 2 = 6$ 
  - (i) truncated to three decimal places
  - (ii) rounded off to three decimal places
2. (a) Using the bisection method, find the root of the equation  $x^3 - x^2 - 1 = 0$ , correct to three decimal places. 7
- (b) Find the missing term in the table using finite difference table 7

$x$	2	3	4	5	6
$y$	45.0	49.2	54.1	...	67.4

**UNIT-II**

4. (a) Apply Gauss-Seidel iteration method to solve the equation 8
  - $20x + y - 2z = 17$
  - $3x + 20y - z = -18$
  - $2x - 3y + 20z = 25$
- (b) Find the eigenvalues and eigenvectors of the matrix

$$A = \begin{bmatrix} 1 & 4 \\ 3 & 2 \end{bmatrix}$$

6

3. (a) What are the pitfalls of Gauss elimination method? Solve the following set of linear equations by Gauss-Jordan method

$$3x + 6y + z = 16 \qquad 2+8=10$$

$$2x + 4y + 3z = 13$$

$$x + 3y + 2z = 9$$

- (b) Convert the following set of equations into matrix form and carry out the partial pivoting 4

$$-3a + 5b + 6c + 2d = 11$$

$$15a + 7b + 6c - 4d = 65$$

$$8a + 12b + 20c + 6 = 60$$

$$3a + 4b + 7c + 9d = 7$$

### UNIT-III

5. (a) For the following data, prepare a forward difference table and using Newton's forward difference formula, find the value of  $y$  for  $x = 8.2$  7

$x$	2	3	4	5	6	7	8	9
$y$	-14	4	40	100	190	316	484	700

- (b) The velocity  $u$  of water flowing over flat surface is measured at several distances  $y$  away from the surface. Fit the curve  $u = ay^2 + by + c$ , using the least squares criterion. 7

$y(\text{cm})$	0	1	2	3	4
$u(\text{cm/s})$	0	5	15	3	50

6. (a) Find the cubic polynomial which takes the following values and evaluate  $f(4)$  7

$x$	0	1	2	3
$f(x)$	1	2	1	10

- (b) Fit a straight line to following data using least square fit method 7

$x$	6	7	7	8	8	8	9	9	10
$y$	5	5	4	5	4	3	4	3	3

## UNIT-IV

7. (a) Evaluate the value of integration  $I = \int_{\frac{\pi}{4}}^{\frac{3\pi}{8}} (\sin x) dx$ . Using the trapezoidal rule with
- (i) stripe width =  $\frac{\pi}{32}$
- (ii) number of stripes = 6
- Compare the answer with analytical answer and determine the error. 6+2+2=10
- (b) Estimate the integration  $f(x) = \int_{2.2}^{3.4} (x^3 - 2x^2 + 7x - 5) dx$  using Simpson's 1/3 rule assuming 8 stripes. 4

8. (a) Using Newton's forward difference formula, find  $\frac{dy}{dx}$  and  $\frac{d^2y}{dx^2}$  at  $x = 1.1$  for the following data 6+2=8

$x$	1	1.1	1.2	1.3	1.4	1.5	1.6
$y$	7.989	8.403	8.781	9.129	9.451	9.75	10.031

- (b) Evaluate  $\int_{-1}^{+1} \frac{dx}{1+x^2}$  using Gauss formula for  $n = 2$  and  $n = 3$ . 3+3+6

## UNIT-V

9. (a) Solve  $\frac{dy}{dx} = 3x + \frac{y}{2}$  with  $y(0) = 1$ ,  $h = 0.1$ , using the 2nd order Runge-Kutta method. 5
- (b) Solve  $\frac{dy}{dx} = \frac{y-x}{y+x}$  using Euler's method to find  $y$  at  $x = 0.1$ .  
Given,  $y(0) = 1$  and  $h = 0.02$ . 9

10. Solve the Laplace's equation  $u_{xx} + u_{yy} = 0$  for the square mesh with boundary values as shown by Jacobi's method. 14

